

Legend to slides-Charles Gerday- Punta del Este-6th May 2010

Slide 1: Title: Psychrophiles: a challenge for life

Conference given by Charles Gerday, Honorary Professor of Biochemistry-University of Liege, Belgium. e-mail: ch.gerday@ulg.ac.be

Slide 2: The Arrhenius law is an equation telling us that the rates of chemical reactions (k) that occur in living organisms, are exponentially depending on temperature (T). This can also be expressed by the Q_{10} , which is the ratio between the rates of a chemical reaction run at two temperatures differing by an interval of 10°C. In living organisms, this Q_{10} is usually between 2-3.

Slide 3: So the question is: "How do psychrophiles maintain appropriate rate of reactions?"

Slide 4: In extremely low temperature environments, below 0°C, the fluids of a living organism are exposed to freezing, so an other question can be addressed: "How do psychrophiles, living below 0°C, prevent freezing?"

Slide 5: This slide shows the effect of freezing of external fluids on unprotected and protected cells. On the left side: a normal cell. On the right side: **below**, an unprotected cell surrounded by external ice crystals. As only pure water freezes there is a rapid and severe concentration of the solutes present outside the cell. This induces the loss of water molecules from the intracellular space, aiming to re-equilibrate the osmotic pressure of both sides of the cell, but leading to cell dehydration and death. **Above**, a protected cell producing in particular cryoprotectants that actively participate in the re-equilibration of the osmotic pressures and prevent the dehydration of the cell.

Slide 6: In sea ice, even at very low temperature down to -20°C, there is a persistence of liquid veins (brine veins), often highly salted, that are populated by numerous bacteria displaying metabolic activities.

Slide 7: This cartoon illustrates the chemical composition of a membrane formed by two layers of lipids which have to be modified to secure an appropriate proton permeability at the respective environmental temperatures of living organisms.

Slide 8: The cold-adapted enzymes produced by psychrophilic organisms have two general properties: a high specific activity at low and moderate temperatures and a high thermosensitivity, two properties that can be extremely valuable for many biotechnological applications.

Slide 9: This represents the relationship between the specific activity of an enzyme (alpha-amylase, hydrolyzing starch) and the temperature of the environment and this for a cold-adapted enzyme (blue curve), and a temperate or mesophilic counterpart (green curve). One can clearly see that the specific activity of the psychrophilic enzyme is much higher than that of the temperate counterpart at low and moderate temperatures. The apparent maximum is shifted towards low temperatures whereas the cold-adapted enzyme is rapidly inactivated at temperature higher than 30°C.

Slide 10: An Antarctic bacterial strain was grown at different temperatures and the graph, on the left, shows that the cell density is much higher at 4°C when compared to 18°C. At 25°C the production of cells is nearly completely abolished as well as that of exogenous enzymes such as the alpha-amylase necessary for the survival of the strain (graph on the right). This clearly shows that cold-adapted cells are very sensitive to heat. They are on the contrary well adapted to their low temperature environment.

Slide 11: To resist freezing, cold-adapted organisms produce antifreeze molecules, often proteins, which prevent the formation of large crystals of ice. The first to be described in the sixties by Art DeVries and co-workers is a glycoprotein isolated from an Antarctic fish and formed by a repetitive sequence of amino acids: Ala-Ala-Thr to which is bound a disaccharide.

Slide 12: Other antifreeze proteins have been discovered in a large variety of organisms. For example, five different families of protein antifreeze have been identified in various fish species adapted to low temperatures. A particularity is that they are not evolutionary related, their structure is different from each other but the effect is similar, although strongly dependent on the respective concentrations. They bind to the micro ice crystals at specific sites and inhibit crystal growing

Slide 13: The re-crystallization of ice is illustrated as well as its inhibition by antifreeze molecules. The ice in the upper panel contains **AFP** (Antifreeze proteins) and crystals are tiny whereas, on the lower panel **AFP's** are not present and the ice crystals

increase in size whereas the number of crystals goes down. (Courtesy of Professor Peter L Davies- Queens University-Canada)

Slide 14: The structure of some cryoprotectants is shown. They contribute to depress the freezing point of physiological fluids and to protect proteins against cold-denaturation. Most of them are monomeric sugars, such as glucose or derivated polyols such as sorbitol.

Slide 15: There are three main effects of cryoprotectants:

- They depress the freezing point of water by colligative effect
- They prevent the dehydration of the cell when extracellular ice is present
- They protect proteins against cold denaturation.

Slide 16: The various effects of ice-structuring proteins are illustrated. In the absence of nucleation pure water freezes around -39°C. Usually however pure water is not that pure and contains tiny particles that favor the nucleation so that ordinary water freezes at 0°C. Some psychrophiles produce, **INP**, for ice nucleating proteins that increase the freezing point of water; they are used in artificial snow production on ski tracks. They can also produce **ANP**, for anti nucleating proteins, which, on the contrary, favor super-cooling, meaning a decrease of the freezing point of water. Ice structuring proteins also include the antifreeze molecules: some (**AFP**) depress the freezing point and prevent the growth of ice crystals, some other (**IR** for ice recrystallization) do not depress the freezing point but prevent the growth of ice crystals in physiological fluids.

Slide 17: Ice nucleating proteins have four main effects:

- Ice is confined into the extracellular space
- The formation of ice is controlled by the organism and progressively formed enabling appropriate metabolic adjustments.
- In this way cryoprotectants can also be produced concomitantly preventing excessive dehydration
- The formation of extracellular ice induces limited cell dehydration and that also contributes to depress the freezing point of the intracellular fluid

Slide 18: Identical to slide **16** in order to comment on **ANP** (Anti-nucleating proteins) that induces super-cooling, meaning the depression of the freezing point of fluids. **ANP** were successfully used in the non-freezing preservation of organs such as livers

at temperatures around -3°C .

Slide 19: The slide illustrates the effect of a psychrophilic bacterial strain (*Arthrobacter psychrolactophilus* isolated near the scientific station of NY-Alesund in Spitzberg) on the de-pollution, at 10°C , of waters contaminated by polymers such as proteins, lipids and sugars. This biotechnological process is named **bio-augmentation** and consists in the addition to the environment of exogenous microorganisms that complete the de-polluting effect of indigenous microorganisms.

Slide 20: This slide illustrates the various biotechnological applications in which psychrophilic microorganisms and their enzymes can actually (or in a near future) play a crucial role. The main fields are: the detergent market, the food industry, organic synthesis, molecular biology, textiles and environment.

Slide 21: A cold-adapted beta-galactosidase has been isolated in our laboratory from an Antarctic strain *Pseudoalteromonas haloplanktis*, in view to be used in the industrial process of lactose hydrolysis in milk. Indeed about $\frac{2}{3}$ of the world population are intolerant to lactose and so dairy products have to be free of lactose. A patent has been taken out by our University and the process is going to be used by our industrial Belgian partner, Nutrilab. The slide shows that the common sugar present in milk, the lactose, responsible for the intolerance to milk, can be enzymatically split into its monomeric constituents, the non-harmful sugars, glucose and galactose, by an enzyme named lactase or beta-galactosidase.

Slide 22: This table is a comparison between the efficiency of a cold-adapted beta-galactosidase produced by an Antarctic strain of *Pseudoalteromonas haloplanktis* in reducing the lactose in milk and the homologous and commercial mesophilic (temperate) counterpart. The figures show that at 5°C the cold-adapted enzyme is nearly three times more efficient.

Slide 23: With an industrial partner we are developing, on the basis of the efficiency of the cold-adapted beta-galactosidase in reducing the amount of lactose in milk, a process that will enable the transformation of the D-galactose produced during the hydrolysis into a high added value new sweetener, the Tagatose.

Slide 24: On this slide is shown the three dimensional structure of a cold-adapted xylanase, isolated in our laboratory, from an Antarctic bacterial strain,

Pseudoalteromonas haloplanktis, that belongs to a new family of xylanases, family 8. The structure is compared to the usual other xylanases which belong to family 10 and family 11.

Slide 25: The cold-adapted xylanase was tested in the baking process of so-called Argentinian breads and other bread forms. The table shows that when compared to its mesophilic counterpart, the cold-adapted enzyme dramatically improves the process since similar loaf volume and cut width are obtained with an amount of enzyme lowered by a factor of 100. We have also a patent for this application and the cold-adapted xylanase is now sold all over the world for the baking industry.

Slide 26: Most of the work carried out in our laboratory has been made possible thanks to our participation to summer campaigns organized by the “Institut français de recherches et de technologies polaires” at the Antarctic station J-S Dumont d’Urville in Terre Adélie. We thank this Institution for their continuous support and accommodation of our research fellows at this station and also at the sub-Antarctic station Port-aux Français at the Kerguelen archipelago for nearly 20 years.

Slide 27: Local and international researchers have actively collaborated with us mainly through European Union research contracts. They are not only outstanding collaborators but also friends (Richard Haser from Lyon, Luisa Tutino from Naples, Nick Russell from London, Rick Cavicchioli from Sydney, Rosa Margesin from Innsbruck, Jos van Beeumen from Gent and Philippe Thonart from Gembloux. Some of the work has also been carried out at the Arctic station of Ny-Alesund in Spitzberg thanks to our accommodation at the Italian house run by the CNR-Italy.